AP® Calculus BC
2002 Free-Response Questions
Form B

These materials were produced by Educational Testing Service® (ETS®), which develops and administers the examinations of the Advanced Placement Program for the College Board. The College Board and Educational Testing Service (ETS) are dedicated to the principle of equal opportunity, and their programs, services, and employment policies are guided by that principle.

The College Board is a national nonprofit membership association dedicated to preparing, inspiring, and connecting students to college and opportunity. Founded in 1900, the association is composed of more than 4,200 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three million students and their parents, 22,000 high schools, and 3,500 colleges, through major programs and services in college admission, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT®, the PSAT/NMSQT®, and the Advanced Placement Program® (AP®). The College Board is committed to the principles of equity and excellence, and that commitment is embodied in all of its programs, services, activities, and concerns.

CALCULUS BC
SECTION II, Part A
Time—45 minutes
Number of problems—3

A graphing calculator is required for some problems or parts of problems.

---

1. A particle moves in the xy-plane so that its position at any time t, for $-\pi \le t \le \pi$, is given by $x(t) = \sin(3t)$ and $y(t) = 2t$.

   (a) Sketch the path of the particle in the xy-plane provided. Indicate the direction of motion along the path. **(Note: Use the axes provided in the test booklet.)**



   (b) Find the range of $x(t)$ and the range of $y(t)$.

   (c) Find the smallest positive value of t for which the x-coordinate of the particle is a local maximum. What is the speed of the particle at this time?

   (d) Is the distance traveled by the particle from $t = -\pi$ to $t = \pi$ greater than 5π ? Justify your answer.

GO ON TO THE NEXT PAGE.

2. The number of gallons, $P(t)$, of a pollutant in a lake changes at the rate $P'(t) = 1 - 3e^{-0.2\sqrt{t}}$ gallons per day, where t is measured in days. There are 50 gallons of the pollutant in the lake at time $t = 0$. The lake is considered to be safe when it contains 40 gallons or less of pollutant.

   (a) Is the amount of pollutant increasing at time $t = 9$? Why or why not?

   (b) For what value of t will the number of gallons of pollutant be at its minimum? Justify your answer.

   (c) Is the lake safe when the number of gallons of pollutant is at its minimum? Justify your answer.

   (d) An investigator uses the tangent line approximation to $P(t)$ at $t = 0$ as a model for the amount of pollutant in the lake. At what time t does this model predict that the lake becomes safe?



3. Let R be the region in the first quadrant bounded by the y-axis and the graphs of $y = 4x - x^3 + 1$ and $y = \frac{3}{4}x$.

   (a) Find the area of R.

   (b) Find the volume of the solid generated when R is revolved about the x-axis.

   (c) Write an expression involving one or more integrals that gives the perimeter of R. Do not evaluate.

END OF PART A OF SECTION II

CALCULUS BC
SECTION II, Part B
Time—45 minutes
Number of problems—3

No calculator is allowed for these problems.



Graph of f

4. The graph of a differentiable function f on the closed interval $[-3, 15]$ is shown in the figure above. The graph of f has a horizontal tangent line at $x = 6$. Let $g(x) = 5 + \int_6^x f(t)\,dt$ for $-3 \le x \le 15$.

   (a) Find $g(6)$, $g'(6)$, and $g''(6)$.

   (b) On what intervals is g decreasing? Justify your answer.

   (c) On what intervals is the graph of g concave down? Justify your answer.

   (d) Find a trapezoidal approximation of $\int_{-3}^{15} f(t)\,dt$ using six subintervals of length $\Delta t = 3$.

---

5. Consider the differential equation $\frac{dy}{dx} = \frac{3 - x}{y}$.

   (a) Let $y = f(x)$ be the particular solution to the given differential equation for $1 < x < 5$ such that the line $y = -2$ is tangent to the graph of f. Find the x-coordinate of the point of tangency, and determine whether f has a local maximum, local minimum, or neither at this point. Justify your answer.

   (b) Let $y = g(x)$ be the particular solution to the given differential equation for $-2 < x < 8$, with the initial condition $g(6) = -4$. Find $y = g(x)$.

GO ON TO THE NEXT PAGE.

6. The Maclaurin series for $\ln\left(\frac{1}{1-x}\right)$ is $\sum_{n=1}^{\infty}\frac{x^n}{n}$ with interval of convergence $-1 \le x < 1$.

   (a) Find the Maclaurin series for $\ln\left(\frac{1}{1+3x}\right)$ and determine the interval of convergence.

   (b) Find the value of $\sum_{n=1}^{\infty}\frac{(-1)^n}{n}$.

   (c) Give a value of p such that $\sum_{n=1}^{\infty}\frac{(-1)^n}{n^p}$ converges, but $\sum_{n=1}^{\infty}\frac{1}{n^{2p}}$ diverges. Give reasons why your value of p is correct.

   (d) Give a value of p such that $\sum_{n=1}^{\infty}\frac{1}{n^p}$ diverges, but $\sum_{n=1}^{\infty}\frac{1}{n^{2p}}$ converges. Give reasons why your value of p is correct.

END OF EXAMINATION